

Semantics for the λ -calculus

Arnoud van der Leer
Delft University of Technology
arnoudvanderleer@gmail.com

2023-11-02

Classical lambda calculus in modern dress

- Paper by Martin Hyland.
- About models for the λ -calculus.
- Three 'big' theorems.
- My job: 'annotate'.

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



The **untyped** λ -calculus

Describes a collection consisting of (only) functions.

The **untyped** λ -calculus

Describes a collection consisting of (only) functions.

Has terms, consisting of *variables*, *application* and *abstraction*:

$$x_1$$
$$x_1(x_2 x_1)$$
$$\lambda x_1, x_1$$
$$\lambda x_3 x_2 x_1, x_1(x_2 x_3).$$

Can have β - and η -equality:

$$(\lambda x_n, f)g = f[x_n := g] \quad \lambda x_n, (fx_n) = f.$$

The **untyped** λ -calculus

Describes a collection consisting of (only) functions.

Has terms, consisting of *variables*, *application* and *abstraction*:

$$\begin{aligned} & x_1 \\ & x_1(x_2 x_1) \\ & \lambda x_1, x_1 \\ & \lambda x_3 x_2 x_1, x_1(x_2 x_3). \end{aligned}$$

Can have β - and η -equality:

$$(\lambda x_n, f)g = f[x_n := g] \quad \lambda x_n, (fx_n) = f.$$

The (*pure*) λ -calculus: Described exactly by the above.

Algebraic theories: objects with variables and substitution

Example

λ -calculus: $\Lambda_n = \{(\lambda x_1, x_1), x_5, (\lambda x_3, x_7)x_{42}\}$.

Algebraic theories: objects with variables and substitution

Example

λ -calculus: $\Lambda_n = \{(\lambda x_1, x_1), x_5, (\lambda x_3, x_7)x_{42}\}$.

Example

Polynomial ring: $\mathbb{Z}[x_1, \dots, x_n] = \{1, x_3, 2048 + 7x_1^{37} - x_6x_{13}^{42}x_{17}^{1729}, \dots\}$.

Algebraic theories: objects with variables and substitution

Example

λ -calculus: $\Lambda_n = \{(\lambda x_1, x_1), x_5, (\lambda x_3, x_7)x_{42}\}$.

Example

Polynomial ring: $\mathbb{Z}[x_1, \dots, x_n] = \{1, x_3, 2048 + 7x_1^{37} - x_6x_{13}^{42}x_{17}^{1729}, \dots\}$.

Definition

An *algebraic theory* T is a sequence of sets T_n with variables $x_{i,n} \in T_n$ (for $0 \leq i < n$) and a substitution operation $\bullet : T_m \times T_n^m \rightarrow T_n$.

λ -theory: structure with app and abs

Definition

A λ -theory L is an algebraic theory, together with abstraction functions $\lambda : L_{n+1} \rightarrow L_n$ and application functions $\rho : L_n \rightarrow L_{n+1}$ (both compatible with the substitution).

λ -theory: structure with app and abs

Definition

A λ -theory L is an algebraic theory, together with abstraction functions $\lambda : L_{n+1} \rightarrow L_n$ and application functions $\rho : L_n \rightarrow L_{n+1}$ (both compatible with the substitution).

The pure λ -calculus Λ is the initial λ -theory.

λ -theory: structure with app and abs

Definition

A λ -theory L is an algebraic theory, together with abstraction functions $\lambda : L_{n+1} \rightarrow L_n$ and application functions $\rho : L_n \rightarrow L_{n+1}$ (both compatible with the substitution).

The pure λ -calculus Λ is the initial λ -theory.

β - and η -equality:

$$\rho_n \circ \lambda_n = \text{Id}_{L_{n+1}} \quad \lambda_n \circ \rho_n = \text{Id}_{L_n}.$$

Algebras: Interpretations (or denotations)

We want to interpret terms with free variables as functions from a context to a set

Example

In $T(n) = \mathbb{Z}[x_1, \dots, x_n]$, we can take a set $A = \mathbb{Q}$ and get

$$2x_1 + 3x_1^2x_2 : A^2 \rightarrow A, \quad (a_1, a_2) \mapsto 2 \cdot a_1 + 3 \cdot a_1^2 \cdot a_2.$$

Algebras: Interpretations (or denotations)

We want to interpret terms with free variables as functions from a context to a set

Example

In $T(n) = \mathbb{Z}[x_1, \dots, x_n]$, we can take a set $A = \mathbb{Q}$ and get

$$2x_1 + 3x_1^2x_2 : A^2 \rightarrow A, \quad (a_1, a_2) \mapsto 2 \cdot a_1 + 3 \cdot a_1^2 \cdot a_2.$$

Definition

For an algebraic theory T , a T -algebra A is a set A , together with interpretation functions $T_n \times A^n \rightarrow A$ for all n (respecting the variables and substitution).

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



Scott's representation theorem (1980)

For every λ -theory L , we can find a category C and an object $X : C_0$, such that L is isomorphic to the endomorphism theory of X : the λ -theory $E(X)$ given by $E(X)_n = X^n \rightarrow X$.

Scott's representation theorem (1980)

For every λ -theory L , we can find a category C and an object $X : C_0$, such that L is isomorphic to the endomorphism theory of X : the λ -theory $E(X)$ given by $E(X)_n = X^n \rightarrow X$.

The variables of $E(X)_n$ are the projections $\pi_i : X^n \rightarrow X$. Also, substituting $g_1, \dots, g_m : X^n \rightarrow X$ into $f : X^m \rightarrow X$ composes f with $\langle g_1, \dots, g_m \rangle : X^n \rightarrow X^m$.

Scott's representation theorem (1980)

For every λ -theory L , we can find a category C and an object $X : C_0$, such that L is isomorphic to the endomorphism theory of X : the λ -theory $E(X)$ given by $E(X)_n = X^n \rightarrow X$.

The variables of $E(X)_n$ are the projections $\pi_i : X^n \rightarrow X$. Also, substituting $g_1, \dots, g_m : X^n \rightarrow X$ into $f : X^m \rightarrow X$ composes f with $\langle g_1, \dots, g_m \rangle : X^n \rightarrow X^m$.

We obtain $\lambda : E(X)_{n+1} \rightarrow E(X)_n$ as

$$\lambda : E(X)_{n+1} = (X^{n+1} \rightarrow X) \simeq (X^n \rightarrow X^X) \xrightarrow{\overline{abs} \circ -} (X^n \rightarrow X) = E(X)_n.$$

for some morphism $\overline{abs} : X^X \rightarrow X$. In the same way, we get $\rho : E(X)_n \rightarrow E(X)_{n+1}$ from a morphism $\overline{app} : X \rightarrow X^X$.

Scott's representation theorem (1980)

For every λ -theory L , we can find a category C and an object $X : C_0$, such that L is isomorphic to the endomorphism theory of X : the λ -theory $E(X)$ given by $E(X)_n = X^n \rightarrow X$.

C is the category of sequences of sets $(P_i)_i$ with a composition $P_m \times L_n^m \rightarrow P_n$ and X is the sequence $(L_i)_i$.

Scott's representation theorem (1980)

For every λ -theory L , we can find a category C and an object $X : C_0$, such that L is isomorphic to the endomorphism theory of X : the λ -theory $E(X)$ given by $E(X)_n = X^n \rightarrow X$.

C is the category of sequences of sets $(P_i)_i$ with a composition $P_m \times L_n^m \rightarrow P_n$ and X is the sequence $(L_i)_i$.

With Hyland's definitions and some lemmas, the representation theorem arises before you know it (*on paper*).

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



“The fundamental theorem of the λ -Calculus”

There is a functor from λ -theories to Λ -algebras, sending L to L_0 : its set of constants.

“The fundamental theorem of the λ -Calculus”

There is a functor from λ -theories to Λ -algebras, sending L to L_0 : its set of constants.

There is also a functor from Λ -algebras to λ -theories. This functor again uses the endomorphism theory $E(X)$ for some object X to construct the λ -theory.

“The fundamental theorem of the λ -Calculus”

There is a functor from λ -theories to Λ -algebras, sending L to L_0 : its set of constants.

There is also a functor from Λ -algebras to λ -theories. This functor again uses the endomorphism theory $E(X)$ for some object X to construct the λ -theory.

Hyland shows that these functors constitute an adjoint equivalence.

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



The category of retracts

Given a λ -theory L , we can view elements $f : L_1$ as one-argument functions, and we can compose them like $f \circ g := f \bullet g$.

Now we construct a category R

$$R_0 = \{a : L_1 \mid a \circ a = a\}, \quad a \rightarrow b = \{f : L_1 \mid b \circ f \circ a = f\}.$$

The category of retracts

Given a λ -theory L , we can view elements $f : L_1$ as one-argument functions, and we can compose them like $f \circ g := f \bullet g$.

Now we construct a category R

$$R_0 = \{a : L_1 \mid a \circ a = a\}, \quad a \rightarrow b = \{f : L_1 \mid b \circ f \circ a = f\}.$$

This category is cartesian closed: it has products, and 'exponentials'. So its morphisms constitute a simply typed λ -calculus: we can do *type theory* with the morphisms.

Relatively cartesian closed

This category is cartesian closed: it has products, and 'exponentials'. So its morphisms constitute a simply typed λ -calculus: we can do *type theory* with the morphisms.

Relatively cartesian closed

This category is cartesian closed: it has products, and 'exponentials'. So its morphisms constitute a simply typed λ -calculus: we can do *type theory* with the morphisms.

If we want to do *dependent type theory*, we need dependent products and sums.

$$\begin{array}{ccc} & \xleftarrow{\Sigma_f} & \\ R/A & \xrightarrow{f^*} & R/B \\ & \xleftarrow{\Pi_f} & \\ \downarrow & & \downarrow \\ A & \xleftarrow{f} & B \end{array}$$

Locally cartesian closed: all pullback functors have both adjoints.

Relatively cartesian closed

This category is cartesian closed: it has products, and 'exponentials'. So its morphisms constitute a simply typed λ -calculus: we can do *type theory* with the morphisms.

If we want to do *dependent type theory*, we need dependent products and sums.

$$\begin{array}{ccc} & \xleftarrow{\Sigma_f} & \\ R/A & \xrightarrow{f^*} & R/B \\ & \xleftarrow{\Pi_f} & \\ \downarrow & & \downarrow \\ A & \xleftarrow{f} & B \end{array}$$

Locally cartesian closed: all pullback functors have both adjoints.

In R , not all pullback functors have both adjoints, but some do: *relatively cartesian closed*.

Relatively cartesian closed

This category is cartesian closed: it has products, and 'exponentials'. So its morphisms constitute a simply typed λ -calculus: we can do *type theory* with the morphisms.

If we want to do *dependent type theory*, we need dependent products and sums.

$$\begin{array}{ccc} & \xleftarrow{\Sigma_f} & \\ R/A & \xrightarrow{f^*} & R/B \\ & \xleftarrow{\Pi_f} & \\ \downarrow & & \downarrow \\ A & \xleftarrow{f} & B \end{array}$$

Locally cartesian closed: all pullback functors have both adjoints.

In R , not all pullback functors have both adjoints, but some do: *relatively cartesian closed*.

I am still working on understanding the proof.

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



Annotating the paper

An algebraic theory T is first a functor $T : \mathbf{F} \rightarrow \mathbf{Sets}$: so we have sets $T(n)$ of n -ary multimaps with variable renamings. In addition, T is equipped with projections $pr_1, \dots, pr_n : T(n)$ including as special case the identity $id \in T(1)$. Finally there are compositions $T(n) \times T(m)^n \rightarrow T(m)$ which are **associative, unital, compatible with projections and natural in n and m** . A map $F : S \rightarrow T$ of algebraic theories is a natural transformation with components $F_n : S(n) \rightarrow T(n)$ preserving projections and composition.

Annotating the paper

An algebraic theory T is first a functor $T : \mathbf{F} \rightarrow \mathbf{Sets}$: so we have sets $T(n)$ of n -ary multimaps with variable renamings. In addition, T is equipped with projections $pr_1, \dots, pr_n : T(n)$ including as special case the identity $id \in T(1)$. Finally there are compositions $T(n) \times T(m)^n \rightarrow T(m)$ which are **associative, unital, compatible with projections and natural in n and m** . A map $F : S \rightarrow T$ of algebraic theories is a natural transformation with components $F_n : S(n) \rightarrow T(n)$ preserving projections and composition.

- Learn the background.
- Decode the definitions and theorems.
- Find examples.
- Formalize (on paper).
- Mechanize.

Mechanization

- Displayed categories:
 - Univalence;
 - Limits (twice);
- Higher inductive types;
- $X^{n+1} = X \times X^n$;
- $X_{n+1} = X_{1+n}$;

Intro

Talking about the λ -calculus

Models

Semantics

The main theorems

Scott's representation theorem

The fundamental theorem of the
 λ -calculus


The category of retracts

My contribution

Annotating the paper

Mechanization

Conclusion



Conclusion

Algebraic theories, λ -theories and their algebras (and 'presheaves') seem to be a promising way to work with models for the λ -calculus.

Conclusion

Algebraic theories, λ -theories and their algebras (and 'presheaves') seem to be a promising way to work with models for the λ -calculus.

3 'big' theorems:

- Every model of the λ -calculus arises as the endomorphism theory of some category.
- There is an equivalence between models of the λ -calculus, and interpretations of the λ -calculus as functions on a set.
- From a model for the untyped λ -calculus, we can create a category in which we can do some form of dependent type theory.

Conclusion

Algebraic theories, λ -theories and their algebras (and 'presheaves') seem to be a promising way to work with models for the λ -calculus.

3 'big' theorems:

- Every model of the λ -calculus arises as the endomorphism theory of some category.
- There is an equivalence between models of the λ -calculus, and interpretations of the λ -calculus as functions on a set.
- From a model for the untyped λ -calculus, we can create a category in which we can do some form of dependent type theory.

I am slowly processing the paper.

Conclusion

Algebraic theories, λ -theories and their algebras (and 'presheaves') seem to be a promising way to work with models for the λ -calculus.

3 'big' theorems:

- Every model of the λ -calculus arises as the endomorphism theory of some category.
- There is an equivalence between models of the λ -calculus, and interpretations of the λ -calculus as functions on a set.
- From a model for the untyped λ -calculus, we can create a category in which we can do some form of dependent type theory.

I am slowly processing the paper.

Mechanization is hard.

Do you have questions?

Do you have questions?

Because I have one: I am still a bit unsure about the exact 'meaning' of relative cartesian closedness. Can someone explain that better to me?